

Title	BLOCH' S CONJECTURE FOR ENRIQUES VARIETIES
Author(s)	Laterveer, Robert
Citation	Osaka Journal of Mathematics. 55(3) p.423-p.438
Issue Date	2018-07
oaire:version	VoR
URL	https://doi.org/10.18910/70080
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

BLOCH'S CONJECTURE FOR ENRIQUES VARIETIES

ROBERT LATERVEER

(Received July 15, 2016, revised January 6, 2017)

Abstract

Enriques varieties have been defined as higher-dimensional generalizations of Enriques surfaces. Bloch's conjecture implies that Enriques varieties should have trivial Chow group of zero-cycles. We prove this is the case for all known examples of irreducible Enriques varieties of index larger than 2. The proof is based on results concerning the Chow motive of generalized Kummer varieties.

1. Introduction

For a smooth complex projective variety X , let $A_j X$ denote the Chow group of dimension j algebraic cycles on X modulo rational equivalence. Let $A_j^{\text{hom}}(X) \subset A_j(X)$ denote the subgroup of homologically trivial cycles. Other than the case of divisors ($j = \dim X - 1$), Chow groups are in general still poorly understood. For example, there is the famous conjecture of Bloch:

Conjecture 1.1 (Bloch [6]). *Let X be a smooth projective complex variety. The following are equivalent:*

- (i) *the Albanese morphism $A_0^{\text{hom}}(X) \rightarrow \text{Alb}(X)$ is an isomorphism;*
- (ii) *the Hodge numbers $h^{j,0}(X)$ are 0 for $j \geq 2$.*

The implication from (i) to (ii) is actually a theorem [21], [8]. The conjectural part is the reverse implication, which has been verified for surfaces of Kodaira dimension less than 2 [7], but is wide open for surfaces of general type (cf. [24], [28] for some examples of surfaces where Conjecture 1.1 is verified).

Interesting examples of varieties with vanishing Hodge numbers $h^{j,0}(X) = 0$ for all $j \geq 1$ are given by *Enriques varieties*. These varieties have been defined and studied by Boissière, Nieper-Wißkirchen and Sarti in [9] (and independently, with a somewhat different definition, by Oguiso–Schröer in [23]). As the name suggests, Enriques varieties are higher-dimensional generalizations of Enriques surfaces. In the same way that Enriques surfaces are closely related to $K3$ surfaces, the study of Enriques varieties is intimately entwined with that of hyperkähler varieties. By definition, an Enriques variety X has the property that some multiple dK_X of the canonical divisor is trivial; the smallest such positive integer d is called the *index* of X .

It is natural to ask whether one can prove Bloch's conjecture for these varieties, i.e.

Conjecture 1.2. *Let X be an Enriques variety (in the sense of [9]). Then*

$$A_0(X) = \mathbb{Z}.$$

The main result of this note gives a partial answer to Conjecture 1.2:

Theorem (=Theorem 3.1). *Let X be an Enriques variety of dimension ≤ 6 . Assume X is a quotient*

$$X = K/G,$$

where $K = K_n(A)$ is a generalized Kummer variety and G is a group of automorphisms acting freely and induced by a finite order automorphism of A . Then

$$A_0(X) = \mathbb{Z}.$$

Theorem 3.1 applies to all known examples of irreducible Enriques varieties with index > 2 (these examples can be found in [9] and [23]). The proof of Theorem 3.1 is a straightforward application of results of Xu [30] and Lin [20], combined with Kimura's theory of *finite-dimensional motives* [19].

As a corollary (Corollary 3.9), varieties as in Theorem 3.1 verify certain cases of the generalized Hodge conjecture.

Conventions 1. *In this note, the word variety will refer to a reduced irreducible scheme of finite type over \mathbb{C} .*

For any variety X , we will denote by $A_j X$ the Chow group of j -dimensional cycles on X , and we will write

$$A_j(X)_{\mathbb{Q}} := A_j(X) \otimes_{\mathbb{Z}} \mathbb{Q}$$

for Chow groups with rational coefficients. For X smooth of dimension n the notations $A_j X$ and $A^{n-j} X$ will be used interchangeably.

The notations $A_{\text{hom}}^j(X)$ and $A_{AJ}^j(X)$ will be used to indicate the subgroups of homologically, resp. Abel–Jacobi trivial cycles. The contravariant category of Chow motives (i.e., pure motives with respect to rational equivalence as in [26], [22]) will be denoted \mathcal{M}_{rat} . The category of pure motives with respect to homological equivalence will be denoted \mathcal{M}_{hom} .

2. Preliminary material

2.1. Quotient varieties.

DEFINITION 2.1. *A projective quotient variety is a variety*

$$X = Y/G,$$

where Y is a smooth projective variety and $G \subset \text{Aut}(Y)$ is a finite group.

Proposition 2.2 (Fulton [12]). *Let X be a projective quotient variety of dimension n . Let $A^*(X)$ denote the operational Chow cohomology ring. The natural map*

$$A^i(X)_{\mathbb{Q}} \rightarrow A_{n-i}(X)_{\mathbb{Q}}$$

is an isomorphism for all i .

Proof. This is [12, Example 17.4.10]. \square

REMARK 2.3. It follows from Proposition 2.2 that the formalism of correspondences goes through unchanged for projective quotient varieties (this is also noted in [12, Example 16.1.13]). We can thus consider motives $(X, p, 0) \in \mathcal{M}_{\text{rat}}$, where X is a projective quotient variety and $p \in A^n(X \times X)_{\mathbb{Q}}$ is a projector. For a projective quotient variety $X = Y/G$, one readily proves (using Manin's identity principle) that there is an isomorphism

$$h(X) \cong h(Y)^G := (Y, \Delta_Y^G, 0) \text{ in } \mathcal{M}_{\text{rat}},$$

where Δ_Y^G denotes the idempotent $\frac{1}{|G|} \sum_{g \in G} \Gamma_g$.

2.2. Finite-dimensionality. We refer to [19], [2], [22], [16] for basics on the notion of finite-dimensional motive. An essential property of varieties with finite-dimensional motive is embodied by the nilpotence theorem:

Theorem 2.4 (Kimura [19]). *Let X be a smooth projective variety of dimension n with finite-dimensional motive. Let $\Gamma \in A^n(X \times X)_{\mathbb{Q}}$ be a correspondence which is numerically trivial. Then there is $N \in \mathbb{N}$ such that*

$$\Gamma^{\circ N} = 0 \quad \in A^n(X \times X)_{\mathbb{Q}}.$$

Actually, the nilpotence property (for all powers of X) could serve as an alternative definition of finite-dimensional motive, as shown by a result of Jannsen [16, Corollary 3.9]. Conjecturally, all smooth projective varieties have finite-dimensional motive [19]. We are still far from knowing this, but at least there are quite a few non-trivial examples:

REMARK 2.5. It is an embarrassing fact that up till now, all examples of finite-dimensional motives happen to lie in the tensor subcategory generated by Chow motives of curves, i.e. they are “motives of abelian type” in the sense of [27]. On the other hand, there exist many motives that lie outside this subcategory, e.g. the motive of a very general quintic hypersurface in \mathbb{P}^3 [11, 7.6].

The notion of finite-dimensionality is easily extended to quotient varieties:

DEFINITION 2.6. Let $X = Y/G$ be a projective quotient variety. We say that X has finite-dimensional motive if the motive

$$h(Y)^G := (Y, \Delta_Y^G, 0) \in \mathcal{M}_{\text{rat}}$$

is finite-dimensional. (Here, Δ_Y^G denotes the idempotent $\frac{1}{|G|} \sum_{g \in G} \Gamma_g \in A^n(Y \times Y)$.)

Clearly, if Y has finite-dimensional motive then also $X = Y/G$ has finite-dimensional motive. The nilpotence theorem extends to this set-up:

Proposition 2.7. *Let $X = Y/G$ be a projective quotient variety of dimension n , and assume X has finite-dimensional motive. Let $\Gamma \in A_{\text{num}}^n(X \times X)_{\mathbb{Q}}$. Then there is $N \in \mathbb{N}$ such that*

$$\Gamma^{\circ N} = 0 \quad \in A^n(X \times X)_{\mathbb{Q}}.$$

Proof. Let $p: Y \rightarrow X$ denote the quotient morphism. We associate to Γ a correspondence $\Gamma_Y \in A^n(Y \times Y)_{\mathbb{Q}}$ defined as

$$\Gamma_Y := {}^t\Gamma_p \circ \Gamma \circ \Gamma_p \in A^n(Y \times Y)_{\mathbb{Q}}.$$

By Lieberman's lemma [27, Lemma 3.3], there is equality

$$\Gamma_Y = (p \times p)^*\Gamma \text{ in } A^n(Y \times Y)_{\mathbb{Q}},$$

and so Γ_Y is $G \times G$ -invariant:

$$\Delta_Y^G \circ \Gamma_Y \circ \Delta_Y^G = \Gamma_Y \text{ in } A^n(Y \times Y)_{\mathbb{Q}}.$$

This implies that

$$\Gamma_Y \in \Delta_Y^G \circ A^n(Y \times Y)_{\mathbb{Q}} \circ \Delta_Y^G,$$

and so

$$\Gamma_Y \in \text{End}_{\mathcal{M}_{\text{rat}}}(h(Y)^G).$$

Since clearly Γ_Y is numerically trivial, and $h(Y)^G$ is finite-dimensional (by assumption), there exists $N \in \mathbb{N}$ such that

$$(\Gamma_Y)^{\circ N} = {}^t\Gamma_p \circ \Gamma \circ \Gamma_p \circ {}^t\Gamma_p \circ \cdots \circ \Gamma_p = 0 \text{ in } A^n(Y \times Y)_{\mathbb{Q}}.$$

Using the relation $\Gamma_p \circ {}^t\Gamma_p = d\Delta_X$, this boils down to

$$d^{N-1} {}^t\Gamma_p \circ \Gamma^{\circ N} \circ \Gamma_p = 0 \text{ in } A^n(Y \times Y)_{\mathbb{Q}}.$$

From this, we deduce that also

$$\Gamma^{\circ N} = \frac{1}{d^{N+1}} \Gamma_p \circ (d^{N-1} {}^t\Gamma_p \circ \Gamma^{\circ N} \circ \Gamma_p) \circ {}^t\Gamma_p = 0 \text{ in } A^n(X \times X)_{\mathbb{Q}}.$$

□

2.3. Enriques varieties.

DEFINITION 2.8 [9]. A smooth projective variety is called *Enriques variety* if the following hold:

- (i) the holomorphic Euler characteristic $\chi(X, \mathcal{O}_X) = 1$;
- (ii) there exists an integer $d \geq 2$ (called the *index* of X) such that the canonical bundle K_X has order d in the Picard group of X , and the fundamental group $\pi_1(X)$ is cyclic of order d .

DEFINITION 2.9 [9]. An Enriques variety is called *irreducible* if the holonomy representation of its universal cover is irreducible.

Theorem 2.10 ([9]). *Let X be an irreducible Enriques variety of index > 2 . Then X is the quotient of an irreducible symplectic holomorphic manifold by a group of automorphisms acting freely.*

Proposition 2.11 ([9]). *There exist irreducible Enriques varieties of dimension 4 and index 3, and of dimension 6 and index 4.*

Proof. This is [9, Proposition 4.1], the idea of which is as follows. Let A be the product of 2 elliptic curves, and let ϕ be a finite order automorphism of A . Consider the generalized Kummer variety $K = K_n(A)$ for $n = 3, 4$. For an appropriate choice of ϕ , the induced automorphism $\psi \in \text{Aut}(K)$ is such that the action on K is free, and the quotient

$$X = K / \langle \psi \rangle$$

is an Enriques variety. □

REMARK 2.12. To the best of my knowledge, there are as yet no examples of Enriques varieties with index > 4 .

REMARK 2.13. In [23], there is a definition of “Enriques manifold” which is a priori slightly different from the definition of Enriques variety. (In [18, Remark 1.3(a)], it is explained there might potentially exist Enriques varieties that are *not* Enriques manifolds.) However, the examples given in Proposition 2.11 are also Enriques manifolds (and actually, these examples are also to be found in [23]).

2.4. Generalized Kummer varieties.

DEFINITION 2.14. Let A be an abelian surface. For any $n \in \mathbb{N}$, let

$$\pi: A^{[n]} \rightarrow A^{(n)}$$

denote the Hilbert–Chow morphism from the Hilbert scheme $A^{[n]}$ to the symmetric product $A^{(n)}$. Let $\sigma: A^{(n)} \rightarrow A$ denote the addition morphism. Consider the composition

$$s: A^{[n]} \xrightarrow{\pi} A^{(n)} \xrightarrow{\sigma} A.$$

The generalized Kummer variety is defined as the fibre

$$K_n(A) := s^{-1}(0).$$

$K_n(A)$ is a hyperkähler variety of dimension $2n - 2$.

DEFINITION 2.15 [9]. An automorphism $\psi \in \text{Aut}(K_n(A))$ is *natural* if ψ is induced by an automorphism of A . More precisely, let $A[n]$ denote the n -torsion points of A , and let $\text{Aut}_{\mathbb{Z}}(A)$ denote the group automorphisms of A . As explained in [9, Section 3.1], there is a well-defined homomorphism

$$A[n] \rtimes \text{Aut}_{\mathbb{Z}}(A) \rightarrow \text{Aut}(K_n(A)).$$

The group of natural automorphisms of $K_n(A)$ is defined as the image of this homomorphism.

Theorem 2.16 (Boissière–Nieper-Wißkirchen–Sarti [9]). *Let $n \geq 3$. Let E denote the exceptional divisor of the birational morphism (obtained from π by restriction)*

$$\pi|_{K_n(A)}: K_n(A) \rightarrow K^{(n)} := \sigma^{-1}(0).$$

An automorphism $\psi \in \text{Aut}(K_n(A))$ is natural if and only if $\psi(E) = E$.

Proof. This is [9, Theorem 3.1]. □

2.5. Motive of a generalized Kummer variety.

NOTATION 2.17. For $n \in \mathbb{N}$, let $P(n)$ be the set of partitions of n . A partition $\lambda \in P(n)$ can be written

$$\lambda = (\lambda_1, \dots, \lambda_{\ell_\lambda}) = 1^{a_1} 2^{a_2} \dots r^{a_r},$$

where ℓ_λ is the length of λ . We define $e(\lambda) := \gcd\{\lambda_1, \dots, \lambda_{\ell_\lambda}\}$.

For any $\lambda \in P(n)$, we write

$$A^{(\lambda)} := A^{(a_1)} \times A^{(a_2)} \times \dots \times A^{(a_r)}.$$

DEFINITION 2.18. A homomorphism of Chow motives $\Gamma: M \rightarrow N$ is *split* if Γ admits a left inverse.

Theorem 2.19 (Xu [30]). *Let $K = K_n(A)$ be a generalized Kummer variety. There is a split homomorphism of Chow motives*

$$\Gamma: h(K) \rightarrow \bigoplus_{\lambda \in P(n)} \bigoplus_{\tau \in A[e(\lambda)]} h(A^{(\lambda)})(n - \ell_\lambda - 2) \text{ in } \mathcal{M}_{\text{rat}}.$$

In particular, K has finite-dimensional motive, in the sense of [19] (and even: K has motive of abelian type, in the sense of [27]).

Proof. This follows from [30, Corollary 2.8], which states more precisely that there is an isomorphism

$$h(A \times K) \cong \bigoplus_{\lambda \in P(n)} \bigoplus_{\tau \in A[e(\lambda)]} h(A^{(\lambda)})(n - \ell_\lambda) \text{ in } \mathcal{M}_{\text{rat}}.$$

Theorem 2.19 is obtained by composing with the split homomorphism

$$h(K) \rightarrow h(A \times K) \rightarrow h(A \times K)(-2),$$

where the first arrow is given by projection on the second summand, and the second arrow is given by intersecting with $x \times K$ where $x \in A$. \square

REMARK 2.20. The fact that generalized Kummer varieties have finite-dimensional motive of abelian type (which was first stated explicitly in [30]) seems to have been folklore knowledge for quite some time. Indeed, as noted in [14, Remark 7.10 and §6.1], this fact follows readily from the results of de Cataldo–Migliorini [10].

We mention in passing that L. Fu, in the course of proving the Beauville–Voisin conjecture for generalized Kummer varieties, had previously developed a motivic decomposition for $K_n(A)$ [13]. Parts of the argument of Xu [30] can already be found in [13].

Things simplify if one is only interested in zero-cycles:

Corollary 2.21. *Let $K = K_n(A)$ be a generalized Kummer variety. There is a split injection*

$$\Gamma_*: A_0(K)_{\mathbb{Q}} \rightarrow A_0(A^{(n)})_{\mathbb{Q}}.$$

Proof. This is a consequence of Theorem 2.19; all summands with $\ell_\lambda < n$ vanish for dimension reasons. \square

Theorem 2.22 (Lin [20]). *Let $K = K_n(A)$ be a generalized Kummer variety. There exists a Chow–Künneth decomposition for K , i.e. a set of mutually orthogonal idempotents $\Pi_0^K, \dots, \Pi_{4n-4}^K$ in $A^{2n-2}(K \times K)_{\mathbb{Q}}$ lifting the Künneth components. Moreover, this decomposition satisfies*

$$(\Pi_0^K + \Pi_2^K + \Pi_4^K + \dots + \Pi_{4n-4}^K)_* = \text{id}: A_0(K)_{\mathbb{Q}} \rightarrow A_0(K)_{\mathbb{Q}}.$$

Proof. This is essentially [20, Proposition 4.5]. It follows from Theorem 2.19 that K is motivated by A (in the sense of Arapura [3]). Thus, [3, Lemma 4.2] implies K verifies the standard Lefschetz conjecture, and so in particular the Künneth components of K are algebraic. Finite-dimensionality then gives a Chow–Künneth decomposition [15, Lemma 5.4]. As for the last statement, this follows from the fact that the Beauville filtration on Chow groups of abelian varieties induces a decomposition

$$A_0(K)_{\mathbb{Q}} = \bigoplus_{j=0}^{2n-2} A_0^{(j)}(K)_{\mathbb{Q}}$$

such that

$$A_0^{(j)}(K)_{\mathbb{Q}} = \begin{cases} (\Pi_{4n-4-j}^K)_* A_0(K)_{\mathbb{Q}} & \text{if } j \text{ is even;} \\ 0 & \text{if } j \text{ is odd} \end{cases}$$

[20, Theorem 1.4]. \square

Using the existence of a Chow–Künneth decomposition, Corollary 2.21 can be made more precise:

Corollary 2.23. *Let $K = K_n(A)$ be a generalized Kummer variety. Let Π_j^K (resp. $\Pi_j^{A^{(n)}}$) be any Chow–Künneth decomposition of K (resp. of $A^{(n)}$). For any j , there are split injections*

$$\Gamma_*: (\Pi_j^K)_* A_0(K)_{\mathbb{Q}} \rightarrow (\Pi_{j+4}^{A^{(n)}})_* A_0(A^{(n)})_{\mathbb{Q}}.$$

Proof. Let Ψ denote a left inverse to the homomorphism

$$\Gamma: h(K) \rightarrow N \text{ in } \mathcal{M}_{\text{rat}},$$

where $N \in \mathcal{M}_{\text{rat}}$ is a short-hand for the right-hand side of Theorem 2.19. As a consequence of Theorem 2.19, there are decompositions

$$\Gamma = \Gamma_0 + \Gamma_1: h(K) \rightarrow h(A^{(n)}) \oplus N_1,$$

$$\Psi = (\Psi_0, \Psi_1): h(A^{(n)}) \oplus N_1 \rightarrow h(K) \text{ in } \mathcal{M}_{\text{rat}}$$

satisfying

$$\Psi \circ \Gamma = \Psi_0 \circ \Gamma_0 + \Psi_1 \circ \Gamma_1 = \text{id}: h(K) \rightarrow h(K) \text{ in } \mathcal{M}_{\text{rat}}.$$

Since Γ_0 sends $H^j(K, \mathbb{Q})$ to $H^{j+4}(A^{(n)}, \mathbb{Q})$, there is a homological equivalence

$$L := \Pi_j^K \circ \Psi_0 \circ \Gamma_0 \circ \Pi_j^K = \Pi_j^K \circ \Psi_0 \circ \Pi_{j+4}^{A^{(n)}} \circ \Gamma_0 \circ \Pi_j^K =: R \text{ in } H^{4n-4}(K \times K, \mathbb{Q}).$$

Since K has finite-dimensional motive, this means the difference $L - R$ is nilpotent. Upon developing, this implies

$$L^{\circ N} = Q_1 + Q_2 + \cdots + Q_N \quad \text{in } A^{2n-2}(K \times K)_{\mathbb{Q}},$$

where Q_j is a composition of L and R in which R occurs at least once.

Applying this to 0-cycles, we obtain in particular

$$(L^{\circ N})_* = (Q_1 + \cdots + Q_N)_* : A_0(K)_{\mathbb{Q}} \rightarrow A_0(K)_{\mathbb{Q}}.$$

Now we note that

$$L_* = (\Pi_j^K)_* : A_0(K)_{\mathbb{Q}} \rightarrow A_0(K)_{\mathbb{Q}}$$

thanks to Corollary 2.21. It follows that

$$\begin{aligned} (\Pi_j^K)_* &= ((\Pi_j^K)^{\circ N})_* = (L^{\circ N})_* = (Q_1 + \cdots + Q_N)_* \\ &= ((\text{something}) \circ R \circ \Pi_j^K)_* \\ &= ((\text{something}) \circ \Pi_{j+4}^{A^{(n)}} \circ \Gamma_0 \circ \Pi_j^K)_* : A_0(K)_{\mathbb{Q}} \rightarrow A_0(K)_{\mathbb{Q}}. \end{aligned}$$

It follows that

$$\text{id} = ((\text{something}) \circ \Pi_{j+4}^{A^{(n)}} \circ \Gamma_0)_* : (\Pi_j^K)_* A_0(K)_{\mathbb{Q}} \rightarrow (\Pi_j^K)_* A_0(K)_{\mathbb{Q}}.$$

This proves Corollary 2.23. □

3. Main result

Theorem 3.1. *Let X be an Enriques variety that is a quotient*

$$X = K/G,$$

where $K = K_n(A)$ is a generalized Kummer variety with $n \leq 4$, and G is a group of automorphisms acting freely and induced by a finite order automorphism of A . Then

$$A_0(X) = \mathbb{Z}.$$

Proof. The theorem is true for $n = 2$, so we will suppose from now on that $n \geq 3$. Thanks to Rojtmán [25], we only need to prove that $A_0(X)_{\mathbb{Q}} = \mathbb{Q}$.

Write $G = \langle \psi \rangle$ where $\psi \in \text{Aut}(K)$ is an automorphism (of order $d = \text{index}(X)$) induced by a finite order automorphism

$$\phi \in A[n] \rtimes \text{Aut}_{\mathbb{Z}}(A) \subset \text{Aut}(A).$$

We will write $\phi = t \circ \phi_0$, where t is a translation on A and ϕ_0 is a group automorphism. Let

$$A' := A / \langle \phi_0 \rangle.$$

The surface A' has at most quotient singularities (note that ϕ and ϕ_0 might well have fixpoints even though ψ is fixpoint free). The action of ϕ_0 must be non-symplectic (for otherwise $p_g(X) = 1$), and so $p_g(A') = 0$.

We have seen that the Künneth components of X are algebraic (this follows from Theorem 2.19, or from the results of [10]). Combined with the fact that X has finite-dimensional

motive, this implies [15, Lemma 5.4] that there exists a Chow–Künneth decomposition $\Pi_j^X \in A^{2n-2}(X \times X)_{\mathbb{Q}}$ for X . To prove Theorem 3.1, it suffices to prove

$$(1) \quad (\Pi_j^X)_* A_0^{\text{hom}}(X)_{\mathbb{Q}} = 0 \quad \text{for all } j.$$

The next lemma enables us to change the Chow–Künneth projectors to our convenience; we are not stuck with one particular Chow–Künneth decomposition.

Lemma 3.2. *Let X be a variety with finite-dimensional motive, and such that the Künneth components of X are algebraic. Let Π_j^X and $\hat{\Pi}_j^X$ be two Chow–Künneth decompositions for X . Then for any i and j , there is equivalence*

$$(\Pi_j^X)_* A_i^{\text{hom}}(X)_{\mathbb{Q}} = 0 \Leftrightarrow (\hat{\Pi}_j^X)_* A_i^{\text{hom}}(X)_{\mathbb{Q}} = 0.$$

Proof. This is well-known, and easily proven. For later use, we prove a slightly more general statement. \square

Lemma 3.3. *Let X be as in Lemma 3.2. Let Π_j^X be a Chow–Künneth decomposition, and let $\hat{\pi}_j^X$ be any (not necessarily idempotent, or orthogonal) cycles mapping to the Künneth components $\pi_j \in H^*(X \times X, \mathbb{Q})$. Then for any i and j , we have*

$$(\hat{\pi}_j^X)_* A_i^{\text{hom}}(X)_{\mathbb{Q}} = 0 \Rightarrow (\Pi_j^X)_* A_i^{\text{hom}}(X)_{\mathbb{Q}} = 0.$$

Proof. We have

$$(\Pi_j^X - \hat{\pi}_j^X) = 0 \quad \text{in } H^{2m}(X \times X, \mathbb{Q})$$

(where $m := \dim X$). From Kimura's nilpotence theorem [19], it follows that there exists $N \in \mathbb{N}$ such that

$$(\Pi_j^X - \hat{\pi}_j^X)^{\circ N} = 0 \quad \text{in } A^m(X \times X, \mathbb{Q}).$$

Developing this expression, we obtain

$$\Pi_j^X = (\Pi_j^X)^{\circ N} = P_1 + P_2 + \cdots + P_m \quad \text{in } A^m(X \times X)_{\mathbb{Q}},$$

where each P_j is a composition of correspondences containing at least one copy of $\hat{\pi}_j^X$. But then the right-hand side acts trivially on $A_i^{\text{hom}}(X)_{\mathbb{Q}}$ (by hypothesis), and hence so does the left-hand side. \square

Let us now return to the Enriques variety $X = K/G$, and let us define cycles

$$\hat{\pi}_j^X := \frac{1}{d} \Gamma_p \circ \Pi_j^K \circ {}^t \Gamma_p \in A^n(X \times X)_{\mathbb{Q}},$$

where $p: K \rightarrow X$ is the quotient morphism, and the Π_j^K are as in Theorem 2.22. It follows from Theorem 2.22 that

$$(\hat{\pi}_j^X)_* A_0(X)_{\mathbb{Q}} \subset (\Pi_j^K)_* A_0(K)_{\mathbb{Q}} = 0 \quad \text{for } j \text{ odd}.$$

In view of Lemma 3.3, it follows that the vanishing (1) holds for all odd j .

It remains to establish the vanishing (1) for even j . The next lemma establishes two easy cases of (1):

Lemma 3.4. *Set-up as in Theorem 3.1. Then*

$$(\Pi_j^X)_* A_0^{\text{hom}}(X)_{\mathbb{Q}} = 0 \quad \text{for } j \geq 4n - 6.$$

Proof. (In view of Lemma 3.2, if the lemma is true for one Chow–Künneth decomposition, it is true for *all* Chow–Künneth decompositions.)

The case $j = 4n - 4$ is obvious (indeed, Π_{4n-4}^X is just $X \times x$ for $x \in X$, and so the action factors over $A_0^{\text{hom}}(x)_{\mathbb{Q}} = 0$). As for the second case, we observe that $H^2(X, \mathcal{O}_X) = 0$ so that $H^2(X, \mathbb{Q})$ is algebraic. By hard Lefschetz, $H^{4n-6}(X, \mathbb{Q})$ is also algebraic. This implies that the Künneth component π_{4n-6}^X in cohomology is supported on $D \times C \subset X \times X$, where $D \subset X$ is a (possibly reducible) divisor and $C \subset X$ is a (possibly reducible) curve. The action of π_{4n-6}^X on $A^j(X)_{\mathbb{Q}}$ factors over $A^j(\bar{D})_{\mathbb{Q}}$ (where $\bar{D} \rightarrow D$ is a desingularisation), hence π_{4n-6}^X acts trivially $A_0(X)_{\mathbb{Q}}$. Applying Lemma 3.3, the same holds for any Chow–Künneth projector Π_{4n-6}^X . \square

We now state an equivariant version of Corollaries 2.21 and 2.23:

Proposition 3.5. *Assumptions as in Theorem 3.1.*

(i) *There is a split injection*

$$\Gamma_*: A_0(X)_{\mathbb{Q}} = A_0(K)_{\mathbb{Q}}^G \rightarrow A_0((A')^{(n)})_{\mathbb{Q}}.$$

(ii) *For any j , there are split injections*

$$(\Pi_j^X)_* A_0(X)_{\mathbb{Q}} \rightarrow (\Pi_{j+4}^{(A')^{(n)}})_* A_0((A')^{(n)})_{\mathbb{Q}}$$

(here, Π_j^X and $\Pi_j^{(A')^{(n)}}$ denote Chow–Künneth decompositions of X , resp. of $(A')^{(n)}$).

Proof. To prove this, one needs to delve a bit into the proof of Theorem 2.19, i.e. one needs to understand Xu’s result [30].

By construction of K , there is a commutative diagram (where vertical arrows are closed inclusions)

$$\begin{array}{ccccccc} K = K_n(A) & \rightarrow & K^{(n)}(A) := \sigma^{-1}(0) & \rightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ A^{[n]} & \rightarrow & A^{(n)} & \xrightarrow{\sigma} & A. \end{array}$$

For any $\lambda \in P(n)$, let $K^\lambda := \ker(s_\lambda)$ where

$$\begin{aligned} s_\lambda: A^{a_1} \times \cdots \times A^{a_r} &\rightarrow A, \\ (x_1, \dots, x_{\ell_\lambda}) &\mapsto \sum_{i=1}^{\ell_\lambda} \lambda_i x_i. \end{aligned}$$

We have a stratification

$$K^\lambda = \coprod_{\tau \in A[e(\lambda)]} K_\tau^\lambda,$$

where

$$K_\tau^\lambda := \{(x_1, \dots, x_{\ell_\lambda}) \in A^{\ell_\lambda} \mid \sum_{i=1}^{\ell_\lambda} \frac{\lambda_i}{e(\lambda)} x_i = \tau\}.$$

Let S_a denote the symmetric group on a elements. The action of $S_\lambda := S_{a_1} \times \cdots \times S_{a_r}$ on $A^\lambda := A^{a_1} \times \cdots \times A^{a_r}$ restricts to K^λ (and to the K_τ^λ); the quotient is denoted $K^{(\lambda)}$ (resp. $K_\tau^{(\lambda)}$).

The natural morphism

$$A^{(\lambda)} := A^\lambda / S_\lambda \rightarrow A^{(n)}$$

induces morphisms

$$K_\tau^{(\lambda)} \rightarrow K^{(n)}(A).$$

Then one defines correspondences

$$\hat{\Theta}_\tau^\lambda := (K_\tau^{(\lambda)} \times_{K^{(n)}(A)} K)_{\text{red}} \in A^*(K_\tau^{(\lambda)} \times K)$$

(here $()_{\text{red}}$ means one takes the subvariety with the reduced scheme structure), and

$$\Delta_{\lambda,\tau} := \frac{1}{d_{\lambda,\tau}} \hat{\Theta}_\tau^\lambda \circ {}^t \hat{\Theta}_\tau^\lambda \in A^{2n-2}(K \times K)_\mathbb{Q}$$

(where $d_{\lambda,\tau} \in \mathbb{Q}$ is some constant). One can then prove (using the Beilinson–Bernstein–Deligne decomposition theorem) there is a decomposition

$$\Delta_K = \sum_{\lambda \in P(n)} \sum_{\tau \in A[e(\lambda)]} \Delta_{\lambda,\tau} \text{ in } A^{2n-2}(K \times K)_\mathbb{Q}$$

[30, Lemma 2.5]. This gives rise to an isomorphism of Chow motives

$$(2) \quad h(K) \cong \bigoplus_{\lambda \in P(n)} \bigoplus_{\tau \in A[e(\lambda)]} h(K_\tau^{(\lambda)})(n - \ell_\lambda) \text{ in } \mathcal{M}_{\text{rat}}$$

[30, Theorem 2.7].

Next, one considers the natural morphism

$$\begin{aligned} \varphi: A \times K_\tau^{(\lambda)} &\rightarrow A^{(\lambda)}, \\ (x, z) &\mapsto t_x(z) \end{aligned}$$

(where t_x is the translation by x), and one proves φ induces an isomorphism of Chow motives

$$(3) \quad \Gamma_\varphi: h(A \times K_\tau^{(\lambda)}) \cong h(A^{(\lambda)}) \text{ in } \mathcal{M}_{\text{rat}}$$

[30, Corollary 2.8]. Combining isomorphisms (3) and (2) gives Theorem 2.19.

Consider now the Enriques variety

$$X = K/G,$$

where G is a group acting freely on K and induced by a finite order automorphism ϕ . Since we are only interested in 0-cycles, we only need to consider the one partition of length n , i.e. $\lambda = (1^n)$. We have that

$$K^{((1^n))} = K_0^{((1^n))}$$

is invariant under the automorphism of $A^{(n)}$ induced by ϕ ; we write $K_0^{((1^n))}/G$ for the quotient. Fibre product gives rise to correspondences

$$\hat{\Theta}_0^{(1^n)}(G) := (K_0^{((1^n))}/G \times_{\frac{K^{(n)}(A)}{G}} X)_{\text{red}} \in A^*(K_0^{((1^n))}/G \times X),$$

$$\Delta_{(1^n),0}^G := \frac{1}{d_{(1^n),0}} \hat{\Theta}_0^\lambda(G) \circ {}^t \hat{\Theta}_0^\lambda(G) \in A^{2n-2}(X \times X)_{\mathbb{Q}}.$$

Taking 0-cycles, we get a commutative diagram

$$\begin{array}{ccc} A_0(K)_{\mathbb{Q}} \cong (\Delta_{(1^n),0})_* A_0(K)_{\mathbb{Q}} & \xrightarrow{({}^t \hat{\Theta}_0^{(1^n)})_*} & A_0(K_0^{((1^n))})_{\mathbb{Q}} \\ \uparrow & & \downarrow \\ A_0(X)_{\mathbb{Q}} \cong (\Delta_{(1^n),0}^G)_* A_0(X)_{\mathbb{Q}} & \xrightarrow{({}^t \hat{\Theta}_0^{(1^n)}(G))_*} & A_0(K_0^{((1^n))}/G)_{\mathbb{Q}} \end{array}$$

By (2), the upper horizontal arrow is an isomorphism (with inverse given by $(\hat{\Theta}_0^{(1^n)})_*$). The vertical arrows are split injections. It follows that the lower horizontal arrow is an isomorphism (with inverse given by $(\hat{\Theta}_0^{(1^n)}(G))_*$).

One checks that the morphism $\phi \in \text{Aut}(A)$ induces a morphism

$$\varphi' : A / \langle \phi \rangle \times K_0^{((1^n))}/G \rightarrow (A')^{(n)}.$$

(Indeed, write $\phi = t \circ \phi_0$, where t is a translation on A and ϕ_0 is a group automorphism. It is readily checked that ϕ_0 commutes with φ , i.e. there is a commutative diagram

$$\begin{array}{ccc} A \times K_0^{((1^n))} & \xrightarrow{\varphi} & A^{(n)} \\ \downarrow \phi_0 \times \phi_0^{(n)} & & \downarrow \phi_0^{(n)} \\ A \times K_0^{((1^n))} & \xrightarrow{\varphi} & A^{(n)}, \end{array}$$

where $\phi_0^{(n)}$ is the morphism induced by ϕ_0 . As for the translation $t = t_a$, where $a \in A[n]$, we have a commutative diagram

$$\begin{array}{ccc} A \times K_0^{((1^n))} & \xrightarrow{\varphi} & A^{(n)} \\ \downarrow t_a \times t^{(n)} & & \downarrow \text{id} \\ A \times K_0^{((1^n))} & \xrightarrow{\varphi} & A^{(n)}. \end{array}$$

This proves the existence of φ' .)

Taking 0-cycles, we get a commutative diagram

$$\begin{array}{ccc} A_0(A \times K_0^{((1^n))})_{\mathbb{Q}} & \xrightarrow{\varphi_*} & A_0(A^{(n)})_{\mathbb{Q}} \\ \uparrow & & \uparrow \\ A_0(A / \langle \phi \rangle \times K_0^{((1^n))}/G)_{\mathbb{Q}} & \xrightarrow{(\varphi')_*} & A_0((A')^{(n)})_{\mathbb{Q}}. \end{array}$$

By (3), the upper horizontal arrow is an isomorphism (with inverse given by a multiple of φ^*). The vertical arrows are split injections. It follows that the lower horizontal arrow is an isomorphism. To prove (i) of Proposition 3.5, we consider the composition

$$A_0(X)_{\mathbb{Q}} \xrightarrow{({}^t \hat{\Theta}_0^{(1^n)}(G))_*} A_0(K_0^{((1^n))}/G)_{\mathbb{Q}} \rightarrow A_0(A / \langle \phi \rangle \times K_0^{((1^n))}/G)_{\mathbb{Q}} \xrightarrow{(\varphi')_*} A_0((A')^{(n)})_{\mathbb{Q}},$$

where the first and last arrow are isomorphisms, and the second arrow (defined in the obvious way) is a split injection.

Statement (ii) of Proposition 3.5 is deduced from (i) using finite-dimensionality; this is the same argument as Corollary 2.23. \square

Using Proposition 3.5, we can establish the required vanishing (1) in some further cases:

Lemma 3.6. *Set-up as in Theorem 3.1. Then*

$$(\Pi_j^X)_* A_0^{\text{hom}}(X)_{\mathbb{Q}} = 0 \quad \text{for } j < 3n - 4.$$

Moreover, if $n = 4$ then

$$(\Pi_8^X)_* A_0^{\text{hom}}(X)_{\mathbb{Q}} = 0.$$

Proof. (Again, in view of Lemma 3.2, if the lemma is true for one Chow–Künneth decomposition, it is true for *all* Chow–Künneth decompositions.)

Thanks to Proposition 3.5(ii), it suffices to prove

$$(\Pi_k^{(A')^{(n)}})_* A_0((A')^{(n)})_{\mathbb{Q}} = 0 \quad \text{for all } k < 3n.$$

Let

$$\Delta_{A'} = \Pi_0^{A'} + \Pi_1^{A'} + \Pi_2^{A'} + \Pi_3^{A'} + \Pi_4^{A'} \quad \text{in } A^2(A' \times A')_{\mathbb{Q}}$$

be a Chow–Künneth decomposition for A' . Since $p_g(A') = 0$ and A' has finite-dimensional motive, we may suppose $\Pi_2^{A'}$ is supported on $D \times D$, with $D \subset A'$ a divisor (in other words, the “transcendental part of the motive” of A' is 0, in the language of [17]). Also we may suppose that $\Pi_0^{A'} = x \times A'$ for $x \in A'$ and $\Pi_1^{A'}$ is supported on $D \times A'$, with $D \subset A'$ a divisor (these are general facts, for the Chow–Künneth decomposition of any surface [17]).

As is well-known, the correspondences $\Pi_k^{(A')^{(n)}}$ are induced by correspondences

$$\Pi_k^{(A')^n} := \sum_{k_1 + \dots + k_n = k} \Pi_{k_1}^{A'} \times \dots \times \Pi_{k_n}^{A'} \in A^{2n}((A')^n \times (A')^n),$$

which define an S_n -invariant Chow–Künneth decomposition of $(A')^n$.

There is a commutative diagram

$$\begin{array}{ccc} A_0((A')^n)_{\mathbb{Q}} & \xrightarrow{(\Pi_k^{(A')^n})_*} & A_0((A')^n)_{\mathbb{Q}} \\ \uparrow & & \downarrow \\ A_0((A')^{(n)})_{\mathbb{Q}} & \xrightarrow{(\Pi_k^{(A')^{(n)}})_*} & A_0((A')^{(n)})_{\mathbb{Q}} \end{array}$$

Suppose now $j < 3n$. Then each summand occurring in the definition of $\Pi_k^{(A')^n}$ contains at least one $\Pi_{k_\ell}^{A'}$ with $k_\ell \leq 2$. But this means (by the choice of $\Pi_*^{A'}$ we have made above) that $\Pi_k^{(A')^n}$ is supported on (divisor) $\times (A')^n$ and so acts trivially 0-cycles:

$$(\Pi_k^{(A')^{(n)}})_* A_0((A')^{(n)})_{\mathbb{Q}} = 0 \quad \text{for all } k < 3n.$$

It only remains to treat the case $n = 4$ and $k = 12$. All the summands containing at least one $\Pi_{k_\ell}^{A'}$ with $k_\ell \leq 2$ act trivially on 0-cycles (for the same reason as above). So we may suppose all the k_ℓ are 3, and we need to prove that

$$(\Pi_3^{A'} \times \Pi_3^{A'} \times \Pi_3^{A'} \times \Pi_3^{A'})_* A_0((A')^{(4)})_{\mathbb{Q}} = 0.$$

But there is a natural isomorphism

$$A_0((A')^{(4)})_{\mathbb{Q}} \cong \left(\sum_{\sigma \in S_4} \Gamma_{\sigma} \right)_{*} A_0((A')^4)_{\mathbb{Q}} \subset A_0((A')^4)_{\mathbb{Q}}.$$

One can check that the correspondences $\sum_{\sigma \in S_4} \Gamma_{\sigma}$ and $\Pi_3^{A'} \times \Pi_3^{A'} \times \Pi_3^{A'} \times \Pi_3^{A'}$ commute [19, Lemma 3.4]. It follows that

$$\left(\Pi_3^{A'} \times \Pi_3^{A'} \times \Pi_3^{A'} \times \Pi_3^{A'} \right)_{*} A_0((A')^{(4)})_{\mathbb{Q}} \cong (\text{Sym}^4 \Pi_3^{A'})_{*} A_0((A')^4)_{\mathbb{Q}},$$

where $\text{Sym}^4 \Pi_3^{A'}$ is the projector defining the Chow motive $\text{Sym}^4 h^3(A')$ in the language of [19, Definition 3.5]. The action of the correspondence $\text{Sym}^4 \Pi_3^{A'}$ on cohomology is projection to $\wedge^4 H^3(A', \mathbb{Q})$, which is one-dimensional (since $\dim H^3(A', \mathbb{Q}) = 4$) and consists of Hodge classes:

$$\wedge^4 H^3(A', \mathbb{Q}) \subset H^{12}((A')^4, \mathbb{Q}) \cap F^6,$$

where F^* denotes the Hodge filtration. This implies that

$$\text{Sym}^4 \Pi_3^{A'} \in \left(H^4((A')^4, \mathbb{Q}) \cap F^2 \right) \otimes \left(H^{12}((A')^4, \mathbb{Q}) \cap F^6 \right) \subset H^{16}((A')^4 \times (A')^4, \mathbb{Q}).$$

Next, we note that the Hodge conjecture is known to be true for self-products of abelian surfaces [1, 7.2.2]. This implies the same is true for the quotient variety $(A')^4$. (Indeed, let $p: A^4 \rightarrow (A')^4$ denote the quotient morphism, and assume $a \in H^*((A')^4, \mathbb{Q})$ is a Hodge class. Then $p^*(a)$ is a cycle class. It follows that $p_* p^*(a)$ is a cycle class, and $p_* p^*(a)$ is a multiple of a because of the isomorphism $H^i((A')^4, \mathbb{Q}) \cong H_{8-i}((A')^4, \mathbb{Q})$.)

Using the truth of the Hodge conjecture, we find that there is a cohomological equality

$$\text{Sym}^4 \Pi_3^{A'} = \gamma \quad \text{in } H^{16}((A')^4 \times (A')^4, \mathbb{Q}),$$

where γ is a cycle supported on $V \times W$ for closed subvarieties $V, W \subset (A')^4$ of codimension 2 resp. 6. Since $(A')^4$ has finite-dimensional motive, this implies (Proposition 2.7) there exists $N \in \mathbb{N}$ such that

$$\left(\text{Sym}^4 \Pi_3^{A'} - \gamma \right)^{\circ N} = 0 \quad \text{in } A^8((A')^4 \times (A')^4)_{\mathbb{Q}}.$$

Developing this expression (and noting that $\text{Sym}^4 \Pi_3^{A'}$ is idempotent), this gives a rational equivalence

$$\text{Sym}^4 \Pi_3^{A'} = (\text{Sym}^4 \Pi_3^{A'})^{\circ N} = S_1 + \cdots + S_m \quad \text{in } A^8((A')^4 \times (A')^4)_{\mathbb{Q}},$$

where each S_i is a composition of correspondences in which γ occurs at least once. But γ acts trivially on 0-cycles (for dimension reasons) and so the right-hand side also acts trivially on 0-cycles, and we are done. \square

Theorem 3.1 can now be proven by combining Lemmas 3.4 and 3.6. Indeed, suppose $n = 3$ or $n = 4$. Then all even integers $j \in [0, 4n - 4]$ are either covered by Lemma 3.4 or covered by Lemma 3.6. It follows there is no Chow–Künneth projector Π_j^X acting non-trivially on $A_0^{\text{hom}}(X)_{\mathbb{Q}}$ (i.e., the vanishing (1) is proven), and so this group is trivial. \square

REMARK 3.7. Clearly, Theorem 3.1 applies to the examples furnished by Proposition 2.11.

REMARK 3.8. Note that the assumption on the group G in Theorem 3.1 is more restrictive than just asking that G is a group of natural automorphisms. Also, the dimension hypothesis $n \leq 4$ was merely made for commodity, and is perhaps not really necessary. However, in view of the fact that all known examples of Enriques varieties dominated by generalized Kummer varieties (are given by Proposition 2.11 and so) fit in with these hypotheses, it seems trifling to worry too much about these restrictions.

As a corollary, some cases of the generalized Hodge conjecture are verified:

Corollary 3.9. *Let X be an Enriques variety as in Theorem 3.1. Then $H^j(X, \mathbb{Q})$ is supported on a divisor for all $j > 0$.*

Proof. As is well-known [8], this holds for any variety with trivial Chow group of zero-cycles. \square

One can also say something about codimension 2 cycles:

Corollary 3.10. *Let X be an Enriques variety as in Theorem 3.1. Then $A_{AJ}^2(X)_{\mathbb{Q}} = 0$.*

Proof. Again, this is true for any variety with trivial Chow group of zero-cycles [8]. \square

ACKNOWLEDGEMENTS. The ideas developed in this note came to fruition after the Strasbourg 2014–2015 groupe de travail based on the monograph [29]. Thanks to all the participants of this groupe de travail for the stimulating atmosphere. Thanks to the referee for insightful comments. Many thanks to Yasuyo, Kai and Len for lots of pleasant lunch breaks.

References

- [1] S. Abdulali: *Filtrations on the cohomology of abelian varieties*; in The arithmetic and geometry of algebraic cycles (B. Brent Gordon et alii, eds.), CRM Proceedings and Lecture Notes, American Mathematical Society Providence 2000.
- [2] Y. André: *Motifs de dimension finie (d'après S.-I. Kimura, P. O'Sullivan, ...)*, Séminaire Bourbaki 2003/2004, Astérisque **299** (2005), Exp. No. 929, 115–145.
- [3] D. Arapura: *Motivation for Hodge cycles*, Advances in Math. **207** (2006), 762–781.
- [4] A. Beauville: *Variétés Kähleriennes dont la première classe de Chern est nulle*, J. Differential Geom. **18** (1983), 755–782.
- [5] A. Beauville: *Sur l'anneau de Chow d'une variété abélienne*, Math. Ann. **273** (1986), 647–651.
- [6] S. Bloch: *Lectures on algebraic cycles*, Duke University, Mathematics Department, Durham, N.C., 1980.
- [7] S. Bloch, A. Kas and D. Lieberman: *Zero cycles on surfaces with $p_g = 0$* , Comp. Math. **33** (1976), 135–145.
- [8] S. Bloch and V. Srinivas: *Remarks on correspondences and algebraic cycles*, Amer. J. Math. **105** (1983), 1235–1253.
- [9] S. Boissière, M. Nieper-Wißkirchen and A. Sarti: *Higher dimensional Enriques varieties and automorphisms of generalized Kummer varieties*, J. Math. Pures Appl. **95** (2011), 553–563.
- [10] M. de Cataldo and L. Migliorini: *The Chow motive of semismall resolutions*, Math. Res. Lett. **11** (2004), 151–170.
- [11] P. Deligne: *La conjecture de Weil pour les surfaces K3*, Invent. Math. **15** (1972), 206–226.
- [12] W. Fulton: *Intersection theory*, Ergebnisse der Mathematik, Springer-Verlag, Berlin, 1984.
- [13] L. Fu: *Beauville-Voisin conjecture for generalized Kummer varieties*, Int. Math. Res. Not. IMRN **12** (2015), 3878–3898, 2014.

- [14] L. Fu, Z. Tian and C. Vial: *Motivic hyperkähler resolution conjecture for generalized Kummer varieties*, arXiv:1608.04968.
- [15] U. Jannsen: *Motivic sheaves and filtrations on Chow groups*; in *Motives* (U. Jannsen et al., eds.), Proc. Sympos. Pure Math. **55**, Part 1 (1994).
- [16] U. Jannsen: *On finite-dimensional motives and Murre's conjecture*; in *Algebraic cycles and motives* (J. Nagel and C. Peters, eds.), Cambridge University Press, Cambridge 2007.
- [17] B. Kahn, J. Murre and C. Pedrini: *On the transcendental part of the motive of a surface*; in *Algebraic cycles and motives* (J. Nagel and C. Peters, eds.), Cambridge University Press, Cambridge 2007.
- [18] J. Kim: *Higher dimensional Enriques varieties with even index*, Proc. Amer. Math. Soc. **141** (2013), 3701–3707.
- [19] S. Kimura: *Chow groups are finite dimensional, in some sense*, Math. Ann. **331** (2005), 173–201.
- [20] H. Lin: *On the Chow group of zero-cycles of a generalized Kummer variety*, Adv. Math. **298** (2016), 448–472.
- [21] D. Mumford: *Rational equivalence of 0-cycles on surfaces*, J. Math. Kyoto Univ. **9** (1969), 195–204.
- [22] J. Murre, J. Nagel and C. Peters: *Lectures on the theory of pure motives*, University Lecture Series **61** Amer. Math. Soc., Providence, 2013.
- [23] K. Oguiso and S. Schröer: *Enriques manifolds*, J. Reine Angew. Math. **661** (2011), 215–235.
- [24] C. Pedrini and C. Weibel: *Some surfaces of general type for which Bloch's conjecture holds*, to appear in *Period Domains, Algebraic Cycles, and Arithmetic*, Cambridge Univ. Press, 2015.
- [25] A.A. Rojzman: *The torsion of the group of 0-cycles modulo rational equivalence*, Ann. Math. **111** (1980), 553–569.
- [26] T. Scholl: *Classical motives*; in *Motives* (U. Jannsen et al., eds.), Proc. Sympos. Pure Math. **55**, Part 1 (1994).
- [27] C. Vial: *Remarks on motives of abelian type*, to appear in *Tohoku Math. J.*
- [28] C. Voisin: *Bloch's conjecture for Catanese and Barlow surfaces*, J. Differential Geometry **97** (2014), 149–175.
- [29] C. Voisin: *Chow Rings, Decomposition of the Diagonal, and the Topology of Families*, Princeton University Press, Princeton and Oxford, 2014.
- [30] Z. Xu: *Algebraic cycles on a generalized Kummer variety*, arXiv:1506.04297v1.

Institut de Recherche Mathématique Avancée
 CNRS – Université de Strasbourg
 7 Rue René Descartes
 67084 Strasbourg CEDEX
 France
 e-mail: robert.laterveer@math.unistra.fr